

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
24 August 2000 (24.08.2000)

PCT

(10) International Publication Number  
WO 00/48581 A3

- (51) International Patent Classification<sup>7</sup>: A61K 31/46, 49, D-79713 Bad Säckingen (DE). FÄRBER, Lothar [DE/DE]; Drosselweg 6, D-90562 Heroldsberg (DE).
- (21) International Application Number: PCT/EP00/01269 (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Department, CH-4002 Basel (CH).
- (22) International Filing Date: 16 February 2000 (16.02.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
9903761.6 18 February 1999 (18.02.1999) GB  
9914949.4 25 June 1999 (25.06.1999) GB  
9927877.2 25 November 1999 (25.11.1999) GB
- (71) Applicant (for all designated States except AT, US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).
- (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M.B.H. [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): MÜLLER, Wolfgang [CH/CH]; Im Rehwechsel 30, CH-4102 Binningen (CH). STRATZ, Thomas [DE/DE]; Purkersdorferstrasse
- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— With international search report.
- (88) Date of publication of the international search report:  
11 January 2001
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF 5-HT<sub>2</sub> RECEPTOR ANTAGONISTS FOR TREATING MUSCULOESKELETAL DISEASES

(57) Abstract: The present invention relates to a new use for compounds having 5-HT<sub>2</sub> (serotonin M) receptor antagonist activity, especially tropisetron, for the manufacture of a pharmaceutical composition for the treatment of a non-inflammatory local disease of the musculo-skeletal system, of a local irritation condition of a joint or tendon sheath, or for the local treatment of a local manifestation at the locomotor apparatus of an inflammatory disease except for a crystal induced arthritis and a living pathogen induced inflammatory disease condition as long as the living pathogen is still present.

F5

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/01269

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 A61K31/46 A61P19/00 A61P29/00				
According to International Patent Classification (IPC) or to both national classification and IPC				
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K A61P				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the International search (name of data base and, where practical, search terms used) BIOSIS, EPO-Internal				
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>				
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
X	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; October 1998 (1998-10) FUJII Y ET AL: "Prophylactic antiemetic therapy with granisetron-dexamethasone combination in women undergoing breast surgery." Database accession no: PREV199800515648 XP002149133 abstract & ACTA ANAESTHESIOLOGICA SCANDINAVICA, vol. 42; no. 9, October 1998 (1998-10), pages 1038-1042, ISSN: 0001-5172  -/-	11-14		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.				
<input checked="" type="checkbox"/> Patent family members are listed in annex.				
<b>* Special categories of cited documents:</b> <table border="0"> <tr> <td>           "A" document defining the general state of the art which is not considered to be of particular relevance            "E" earlier document but published on or after the international filing date            "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)            "O" document referring to an oral disclosure, use, exhibition or other means            "P" document published prior to the international filing date but later than the priority date claimed         </td> <td>           "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention            "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone            "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.            "&amp;" document member of the same patent family         </td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family			
Date of the actual completion of the international search  4 October 2000		Date of mailing of the international search report  16/10/2000		
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  Orviz Diaz, P		

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/01269

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE BIOSIS 'Online!  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US;  31 July 1998 (1998-07-31)  GIORDANO JAMES ET AL: "Topical ondansetron  attenuates nociceptive and inflammatory  effects of intradermal capsaicin in  humans."  Database accession no. PREV199800438279  XP002149134  abstract  &amp; EUROPEAN JOURNAL OF PHARMACOLOGY,  vol. 354, no. 1,  31 July 1998 (1998-07-31), pages R13-R14,  ISSN: 0014-2999</p>	1-11,13, 14
X	<p>DATABASE BIOSIS 'Online!  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US; 1996  PRAMANIK S S ET AL: "Serotonin-3 (5HT-3)  receptor antagonists modulate  antinociceptive responses in mice."  Database accession no. PREV199799649054  XP002149135  abstract  &amp; INDIAN JOURNAL OF PHYSIOLOGY AND ALLIED  SCIENCES,  vol. 50, no. 4, 1996, pages 168-172,  ISSN: 0367-8350</p>	1-11,13, 14
X	<p>DATABASE BIOSIS 'Online!  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US; 1996  SAMBORSKI WLODZIMIERZ ET AL: "The 5-HT-3  blockers in the treatment of the primary  fibromyalgia syndrome: A 10-day open study  with tropisetron at a low dose."  Database accession no. PREV199799477436  XP002149136  abstract  &amp; MATERIA MEDICA POLONA,  vol. 28, no. 1, 1996, pages 17-19,  ISSN: 0025-5246</p>	1-11,13, 14

-/-

## INTERNATIONAL SEARCH REPORT

 International Application No  
 PCT/EP 00/01269

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1996 FUJII YOSHITAKA ET AL: "Granisetron and dexamethasone provide more improved prevention of postoperative emesis than granisetron alone in children." Database accession no. PREV199799339428 XP002149137 abstract & CANADIAN JOURNAL OF ANAESTHESIA, vol. 43, no. 12, 1996, pages 1229-1232, ISSN: 0832-610X	11-14
X	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1995 MOSER PAUL C: "The effect of 5-HT-3 receptor antagonists on the writhing response in mice." Database accession no. PREV199598457877 XP002149138 abstract & GENERAL PHARMACOLOGY, vol. 26, no. 6, 1995, pages 1301-1306, ISSN: 0306-3623	1-11,13, 14
X	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1994 SCHWOERER H ET AL: "Treatment of acute gouty arthritis with the 5-hydroxytryptamine antagonist ondansetron." Database accession no. PREV199598040148 XP002149139 abstract & CLINICAL INVESTIGATOR, vol. 72, no. 10, 1994, pages 811-813, ISSN: 0941-0198	1-11,13, 14
X	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1989 GIORDANO J ET AL: "PERIPHERALLY ADMINISTERED SEROTONIN 5-HT-3 RECEPTOR ANTAGONISTS REDUCE INFLAMMATORY PAIN IN RATS" Database accession no. PREV199089030180 XP002149140 abstract & EUROPEAN JOURNAL OF PHARMACOLOGY, vol. 170, no. 1-2, 1989, pages 83-86, ISSN: 0014-2999	1-11,13, 14

-/--

## INTERNATIONAL SEARCH REPORT

Int. onal Application No.

PCT/EP 00/01269

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE BIOSIS 'Online!  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US; 1989  GIORDANO J ET AL: "DIFFERENTIAL ANALGESIC  ACTIONS OF SEROTONIN 5-HT-3 RECEPTOR  ANTAGONISTS IN THE MOUSE"  Database accession no. PREV198988019024  XPO02149141  abstract  &amp; NEUROPHARMACOLOGY,  vol. 28, no. 4, 1989, pages 423-428,  ISSN: 0028-3908</p>	1-11,13, 14
X	<p>DATABASE BIOSIS 'Online!  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US; 1989  ESCHALIER A ET AL: "INFLUENCE OF A  SPECIFIC 5-HT-3 ANTAGONIST ON  CARRAGEENAN-INDUCED HYPERALGESIA IN RATS"  Database accession no. PREV198987128603  XPO02149142  abstract  &amp; PAIN,  vol. 36, no. 2, 1989, pages 249-256,  ISSN: 0304-3959</p>	1-11,13, 14
E	<p>WO 00 48597 A (NOVARTIS ERFIND VERWALT  GMBH ;NOVARTIS AG (CH); MUELLER WOLFGANG  ()) 24 August 2000 (2000-08-24)  the whole document</p>	1-14
X	<p>EP 0 507 637 A (ERBAMONT INC).  7 October 1992 (1992-10-07)  the whole document</p>	1-11,13, 14
X	<p>WO 94 01095 A (SMITHKLINE BEECHAM PLC  ;SANGER GARETH JOHN (GB); BANNER STEPHEN  ED) 20 January 1994 (1994-01-20)  the whole document</p>	1-11,13, 14
X	<p>DOAK G J ET AL: "FORMALIN-INDUCED  NOCICEPTIVE BEHAVIOR AND EDEMA:  INVOLVEMENT OF MULTIPLE PERIPHERAL  5-HYDROXYTRYPTAMINE RECEPTOR SUBTYPES"  NEUROSCIENCE,US,NEW YORK, NY,  vol. 80, no. 3, 1997, pages 939-949,  XP000886556  ISSN: 0306-4522  the whole document</p>	1-11,13, 14
	---	
	-/--	

## INTERNATIONAL SEARCH REPORT

Inter. nat. Application No.

PCT/EP 00/01269

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ROSENSTEIN D M ET AL: "ANTINOCICEPTIVE EFFECTS OF MICRODIALYSIS ADMINISTRATION OF 5-HT1A AND 5-HT RECEPTOR AGONISTS AND ANTAGONISTS IN A MODEL OF ACUTE ARTHRITIS" SOCIETY FOR NEUROSCIENCE ABSTRACTS,US,SOCIETY FOR NEUROSCIENCE, vol. 23, no. 23, 1997, page 1540 XPO00886228 ISSN: 0190-5295 the whole document	1-11,13, 14
X	CUNNINGHAM D ET AL: "OPTIMUM ANTI-EMETIC THERAPY FOR CISPLATIN INDUCED EMESIS OVER REPEAT COURSES: ONDANSETRON PLUS DEXAMETHASONE COMPARED WITH METOCLOPRAMIDE; DEXAMETHASONE PLUS LORAZEPAM" ANNALS OF ONCOLOGY,NL,KLUWER, DORDRECHT, vol. 7, no. 3, 1996, pages 277-282, XPO00886538 ISSN: 0923-7534 the whole document	11-14
X	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1998 BOTELLA A ET AL: "Intracolonic glycerol induces abdominal contractions in rats: Role of 5-HT3 receptors." Database accession no. PREV199900012906 XPO02149157 abstract & FUNDAMENTAL & CLINICAL PHARMACOLOGY, vol. 12, no. 6, 1998, pages 619-623, ISSN: 0767-3981	1-11,13, 14
X	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1998 SAMBORSKI WLODZIMIERZ ET AL: "New concept of pharmacological treatment of fibromyalgia. Do exist two subgroups of fibromyalgia patients?" Database accession no. PREV199800453416 XPO02149158 abstract & REUMATOLOGIA (WARSAW), vol. 36, no. 2, 1998, pages 144-150, ISSN: 0034-6233	1-11,13, 14

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/01269

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0048597	A	24-08-2000	NONE	
EP 0507637	A	07-10-1992	US 5122528 A	16-06-1992
			AU 1402092 A	08-10-1992
			CA 2063912 A	06-10-1992
			JP 6172172 A	21-06-1994
			ZA 9202145 A	27-01-1993
WO 9401095	A	20-01-1994	AU 4507693 A	31-01-1994
			EP 0650355 A	03-05-1995
			JP 7508530 T	21-09-1995



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup>:</b> <b>A61K 31/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/48581</b>  <b>(43) International Publication Date:</b> 24 August 2000 (24.08.00)
<b>(21) International Application Number:</b> PCT/EP00/01269 <b>(22) International Filing Date:</b> 16 February 2000 (16.02.00)  <b>(30) Priority Data:</b> 9903761.6            18 February 1999 (18.02.99)    GB 9914949.4           25 June 1999 (25.06.99)            GB 9927877.2           25 November 1999 (25.11.99)       GB  <b>(71) Applicant (for all designated States except AT US):</b> NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).  <b>(71) Applicant (for AT only):</b> NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M.B.H. [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> MÜLLER, Wolfgang [CH/CH]; Im Rehwechel 30, CH-4102 Binningen (CH). STRATZ, Thomas [DE/DE]; Purkersdorferstrasse 49, D-79713 Bad Säckingen (DE). FÄRBER, Lothar [DE/DE]; Drosselweg 6, D-90562 Heroldsberg (DE).		<b>(74) Agent:</b> BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Department, CH-4002 Basel (CH).  <b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> USE OF 5-HT <sub>3</sub> RECEPTOR ANTAGONISTS  <b>(57) Abstract</b>  The present invention relates to a new use for compounds having 5-HT <sub>3</sub> (serotonin M) receptor antagonist activity, especially tropisetron, for the manufacture of a pharmaceutical composition for the treatment of a non-inflammatory local disease of the musculo-skeletal system, of a local irritation condition of a joint or tendon sheath, or for the local treatment of a local manifestation at the locomotor apparatus of an inflammatory disease except for a crystal induced arthritis and a living pathogen induced inflammatory disease condition as long as the living pathogen is still present.		



*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

### Use of 5-HT<sub>3</sub> Receptor Antagonists

The present invention relates to a new use, in particular a new pharmaceutical use for compounds having 5-HT<sub>3</sub> (serotonin M) receptor, in particular specific 5-HT<sub>3</sub> receptor, antagonist activity, especially in the manufacture of a pharmaceutical composition.

Specifically, the present invention relates to the treatments defined below.

The 5-HT<sub>3</sub>-receptor antagonists comprise a defined and recognised class of pharmaceutically active compounds well known in the art and characterised, as their name implies, by their pharmacological activity. Various 5-HT<sub>3</sub> receptor antagonist compounds are commercially available and clinically applied, e.g. in the treatment of emesis.

In accordance with the present invention it has now surprisingly been found that 5-HT<sub>3</sub> receptor antagonists are useful for the treatment of local non-inflammatory, local irritation-related and local inflammatory disease conditions. This is surprising in that (a) the 5-HT<sub>3</sub> receptor antagonists are effective alone; (b) do not only bring pain relief, but also are effective in the treatment of other symptoms, such as effusion, swelling, stiffness and the like, and (c) are locally effective, thus not having to rely on systemic administration that may show effects e.g. by working via effects on nerves and/or synapses at the spinal cord. It is also astonishing that relatively high local doses are both tolerated and effective when local administration is employed.

Hence, the present invention relates to the use of a 5-HT<sub>3</sub> receptor antagonist or of a pharmaceutically acceptable salt of such an antagonist for the manufacture of a pharmaceutical composition for the treatment of a non-inflammatory local disease of the musculo-skeletal system, of a local irritation condition of a joint or tendon sheath, or for the local treatment a local manifestation at the locomotor apparatus of an inflammatory disease except for a crystal induced arthritis and a living pathogen induced inflammatory disease condition as long as the living pathogen is still present for example the treatment of any process, condition, event, or disease as hereinafter described. In particular, the present invention provides the use as mentioned before where, in addition to pain at least one

further sequela or symptom of the local disease, local irritation condition or local manifestation is alleviated, ameliorated or controlled.

Any 5-HT<sub>3</sub> receptor antagonist can be used in accordance with the invention. Preferred 5-HT<sub>3</sub> receptor antagonists which may be employed in accordance with the present invention are:

- A) Ondansetron [1,2,3,9-tetrahydro-9-methyl-3-[(2-methyl-1H-imidazol-1-yl)methyl]-4H-carbazol-4-one (cf. Merck Index, twelfth edition, item 6979);
- B) Granisetron [endo-1-methyl-N-(9-methyl-9-aza-bicyclo[3.3.1]non-3-yl)-1H-imidazole-3-carboxamide: (cf. loc. cit., item 4557); and
- C) Dolasetron [1H-indole-3-carboxylic acid (2 $\alpha$ , 6 $\alpha$ , 8 $\alpha$ , 9 $\alpha\beta$ )-octahydro-3-oxo-2,6-methano-2H-quinolizin-8-yl ester] (cf. loc. cit., item 3471).

Particular 5-HT<sub>3</sub> receptor antagonists which may be employed in accordance with the present invention are those of the formula 1 as defined in European Patent Publication 189002 B1, the contents of which are incorporated herein by reference, in particular the compound:

- D) Indol-3-yl-carboxylic acid-endo-8-methyl-8-aza-bicyclo[3,2,1]-oct-3-yl-ester, also known as tropisetron: (cf. loc.cit., item 9914).

Further 5-HT<sub>3</sub> receptor antagonists which may be used preferably in accordance with the present invention are:

- E) 4,5,6,7-tetrahydro-5-[(1-methyl-indol-3-yl)carbonyl]benzimidazole (see also ramosetron, see U.S. patent 5,344,927);
- F) (+)-10-methyl-7-(5-methyl-1H-imidazol-4-ylmethyl)-6,7,8,9-tetrahydropyrido[1,2-a]indol-6-one (see also fabesetron, EP 0 361 317); and

- G) [N-(1-ethyl-2-imidazolin-2-yl-methyl)-2-methoxy-4-amino-5-chlorobenzamide (see also lintopride - Chem.- Abstr.-No. 107429-63-0).

A further 5-HT<sub>3</sub> receptor antagonists which may be used preferably in accordance with the present invention is

- (H) 2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one (see also alosetron, EP 0 306 323).

5-HT<sub>3</sub>-receptor antagonists may be employed in accordance with the invention in free or in pharmaceutically acceptable salt form, e.g. as known in the art, for example, in the case of compounds A) to D) above in pharmaceutically acceptable acid addition salt form, for example, in the case of: compound A) the hydrochloride dihydrate; compound B) the hydrochloride; compound C) the mesylate; and compound D) the monohydrochloride. References to 5-HT<sub>3</sub> receptor antagonists collectively or individually throughout the present specification and claims are accordingly to be understood as embracing both free compounds and such pharmaceutically acceptable salt forms, e.g. as clinically employed, and further also solvates, e.g. hydrates, or specific crystal forms of any of these compounds or salts.

For use in accordance with the present invention tropisetron (especially in the formulation called NAVOBAN®) is most preferred.

Thus, the invention relates to the use of a 5-HT<sub>3</sub> receptor antagonist or of a pharmaceutically acceptable salt of such an antagonist for the manufacture of a pharmaceutical composition for the treatment of a non-inflammatory local disease of the musculo-skeletal system, of a local irritation condition of a joint or tendon sheath, or for the local treatment a local manifestation at the locomotor apparatus of an inflammatory disease except for a crystal induced arthritis and a living pathogen induced inflammatory disease condition as long as the living pathogen is still present, where the 5-HT<sub>3</sub> receptor antagonist is selected from the group consisting of ondansetron, granisetron, dolasetron, tropisetron, ramosetron, fabesetron, lintopride and alosetron, which may be used in free form or as a pharmaceutically acceptable salt.

In accordance with the present invention it has now surprisingly been found that 5-HT<sub>3</sub> receptor antagonists are useful for the treatment of certain diseases, processes, conditions or events, namely non-inflammatory local diseases of the musculo-skeletal system; local irritation conditions of the joints or tendon sheaths; or the local treatment also of local manifestations at the locomotor apparatus of inflammatory diseases; such as conditions, processes or events that are due to trauma (including surgery or preferably accident); overload; posture fault; degenerative processes; conditions subsequent to another disease (including infection, for example viral infection, or tumor diseases); or other conditions that result in irritation of the locomotor apparatus of the body; meaning especially therapy of the respective non-inflammatory disease, irritation or manifestation as such, its sequelae or its symptoms, or any combinations of these.

"Treatment" as used herein includes use for the alleviation, amelioration or control of said diseases, processes, conditions or events. It also includes intervention for the alleviation, amelioration or control of the sequelae or symptoms of any one or more of these diseases, for example degeneration (e.g. of cells, epithelia or tissues), swelling, exudation, effusion or pain, stiffness or inflexibility of joints. In this context the term "treatment" is further to be understood as embracing use to reverse, restrict or control progression of any specified disease, process, condition event or the like, including use for disease modifying effect. If any of the mentioned diseases, processes, conditions, manifestations or events, especially an inflammatory disease, process, condition, manifestation or event, is associated with pain, the term "treatment" preferably encompasses the alleviation, amelioration or control (including temporal or permanent removal) of at least one further sequela or symptom in addition to pain, such as swelling, effusion, exsudation, lack of flexibility (e.g. due to stiffness and/or lack of flexibility of joints), loss of strength or degeneration, more preferably of all symptoms and most preferably of the total clinical picture of the respective disease, irritation or manifestation. According to the present invention, the term "treatment" preferably does not have the meaning of prevention.

The present invention is in particular applicable to the treatment of:

(1) non-inflammatory local diseases of the musculo-skeletal system (including a joint or another part of the locomotor apparatus), such as

(1a) myofasciale syndrome, including myelogelosis, chronic lumbalgia or cervicalgia, for example unspecific or in the context of degenerative spinal affections, of static posture fault or malformation of the spine, e.g. cervical syndrome,

(1b) tendomyosis, tendinosis, insertion tendopathy (e.g. epicondylitis), bursopathy or peri-arthropathy,

(1c) overload syndrome of the muscle,

(1d) syndromes due to the compression of nerves or neuropathy (such as medianus compression syndrome = carpal tunnel syndrome), or

(1e) algodystrophy (also called neurodystrophy); or

(2) local irritation conditions (= states of irritation) of a joint or tendon sheath especially related to

(2a) a meniscus lesion (e.g. following surgical intervention or preferably due to damage to the meniscus due to a different cause),

(2b) arthrosis, such as gonarthrosis,

(2c) trauma, including accident or post-operative trauma, e.g. after implant or insertion surgery or endoscopy,

(2d) osteochondritis dissecans, osteonecrosis or joint chondromatosis,

(2e) or various non-inflammatory rheumatoid diseases,

preferably to the local treatment of one or more of the mentioned diseases under (1) and/or

(2);

or

(3) to the local treatment also of a local manifestation at the locomotor apparatus of an inflammatory disease, such as various inflammatory rheumatoid diseases (except for crystal induced arthritis (e.g. gout) and except for living pathogen induced diseases as long as the living pathogen (bacterium, virus or fungus, protozoon, parasite or the like) is still present), such as

(3a) chronic polyarthritis (= rheumatoid arthritis), including juvenile arthritis or psoriasis arthropathy;

(3b) sarcoidosis,

(3c) paraneoplastic syndrome or tumor-induced inflammatory diseases,

(3d) turbid effusions,

(3e) collagenosis, such as systemic Lupus erythematosus, poly-myositis, dermato-myositis, systemic sclerodermia or mixed collagenosis;

(3f) postinfectious arthritis (where no living pathogenic organism can be found at or in the affected part of the body), or

(3g) seronegative spondylarthritis, such as spondylitis ankylosans;

or further

(3h) vasculitis.

In the case of the inflammatory diseases, diseases where a living pathogen, e.g. a virus, a bacterium, a fungus, a protozoon or a parasite or the like, is still present, the treatment of must first aim at removal of the pathogen causative for the disease, before treatment with a 5-HT<sub>3</sub> antagonist is used, as otherwise there is the danger that the causative pathogen remains intact. Then the mere symptomatic treatment with a 5-HT<sub>3</sub> antagonist is contraindicated in order to avoid survival or even further spread of the causative infection. This is also valid in the case of combination with an anti-inflammatory glucocorticosteroid as described in the following, as is the proviso that treatment of crystal-induced inflammation is excluded.

The term "locomotor apparatus" refers to any component of the musculo-skeletal system of the body, especially to bony tissue, muscle, tendons, ligaments, joints, cartilage, perichondrium, periosteum, synovial membrane, bursa and the like.

"Trauma" refers preferably to operative and more preferably to trauma by accident, such as overload, tumble or push.

The term "local treatment" refers to the treatment with one or more 5-HT<sub>3</sub> receptor antagonists near or at the site of the manifestation of the disease to be treated, e.g. by intra-muscular injection, intra-articular injection, or any other injection near or at the site of disease manifestation (that is, preferably the administration has the goal to provide for locally higher concentrations of the administered compound than would be expected to be achieved by systemic administration, or the goal is not systemic exposure), preferably within an area within 20, more preferably 10 cm, still more preferably within 5 cm, around the outer limitation of the manifestation of the local disease, most preferably directly at the affected

are or site, e.g. the are or site of symptom manifestation, such as the area of greatest pain, such as an insertion point, a trigger point or a joint, or also in a broader aspect of the invention by local tissue infiltration or transdermal administration at the site of the manifestation of the disease, e.g. by means of topical administration e.g. by use of gels, creams or ointments or the like, or by transdermal patch technology. In the case of the non-inflammatory diseases mentioned above under (1) and (2), also systemic treatment is possible, e.g. by enteral, especially peroral, e.g. by use of tablets or capsules, or rectal, e.g. by use of enema or suppositories; subcutaneous, intraperitoneal or intra-muscular injection; or infusion is possible. In the case of intravenous administration bolus injection is preferred. However, local treatment is preferred in all cases. An advantage of local treatment is that high efficiency can be reached and that systemic exposure to a 5-HT<sub>3</sub> antagonist can be diminished or avoided. One preferred local way of administration is the intra-articular injection in case of diseases, conditions etc. that relate to joints, e.g. bursopathy or synovitis.

A preferred example of a local manifestation at the locomotor apparatus of an inflammatory disease is synovial inflammation, for example, synovitis, including any of the particular forms of synovitis recited in Dorland's Illustrated Medical Dictionary, 26th edition, pub. W. B. Saunders and Co. at page 1301, as far as it is not crystal-induced. Such synovial inflammation may for example, be consequential to or associated with disease, e.g. arthrosis, including arthritis, e.g. osteoarthritis, rheumatoid arthritis or arthritis deformans.

Further preferred local diseases to be treated according to the invention are those mentioned in the examples.

From the foregoing it will be noticed that the present invention is to be understood especially as embracing the treatment, e.g. therapy, of any disease, process, symptom, event or condition as set forth above, for example for the alleviation or control of non-inflammatory local diseases or irritations or local inflammatory processes or events and the sequelae associated therewith or consequential thereto, e.g. to alleviate or control joint irritation or effusion; with the proviso that if pain is treated, then in addition at least one other disease manifestation (e.g. effusion, swelling, exudation or degeneration) is treated.



Preferably the present invention relates to the treatment, especially the local treatment, of a disease as mentioned above under (2) or (3), more preferably one of the diseases mentioned there other than algodystrophy and vasculitis.

In a further aspect it has been found in accordance with the present invention that 5-HT<sub>3</sub> receptor antagonists are useful as replacement therapy for local glucocorticosteroid, e.g. cortisone or the like, therapy; for example for use in any means of treatment as hereinbefore and hereinafter set forth, e.g. the diseases mentioned under (2) and (3) hereinabove.

The term "replacement therapy" as used herein is to be understood as embracing both use "as full replacement", i.e. use instead of glucocorticosteroid therapy, as well as use "as partial replacement" for glucocorticosteroid therapy, i.e. for administration together with glucocorticosteroid therapy or as a means of reducing glucocorticosteroid dosage or to achieve a glucocorticosteroid sparing effect.

The present invention accordingly provides:

- I. A method of treating any process, condition, event, or disease as hereinbefore set forth, in a subject in need thereof, which method comprises locally administering an effective amount of a 5-HT<sub>3</sub> receptor antagonist, especially locally at or near the site of the local disease, local irritation condition or local manifestation;
- II. A method of providing replacement therapy for glucocorticosteroid therapy in a subject receiving such glucocorticosteroid therapy for or in the treatment of any process, condition, event or disease as hereinbefore set forth, which process comprises locally administering to said subject an effective amount, e.g. a glucocorticosteroid sparing amount, of a 5-HT<sub>3</sub>-receptor antagonist; as well as
- III. A method of treating any process, condition, event or disease as hereinbefore set forth, in a subject in need thereof, which method comprises locally administering an effective amount of a 5-HT<sub>3</sub> receptor antagonist together with a glucocorticosteroid.

Where the term "glucosteroid" is used, this means an anti-inflammatory glucosteroid.

Where co-administration is practiced as under III above the drug substances, i.e. 5-HT<sub>3</sub> receptor antagonist and glucocorticosteroid may be administered sequentially or simultaneously or substantially simultaneously, e.g. employing a fixed combination dosage form.

In further aspects the present invention also provides:

- IV. A 5-HT<sub>3</sub> receptor antagonist for use in, or for use in the manufacture of a pharmaceutical composition for use in; or the use of a pharmaceutical composition comprising a 5-HT<sub>3</sub> receptor antagonist for local use:
- a) in the treatment of any process, condition, event or disease as hereinbefore set forth;
  - b) as replacement therapy for glucocorticosteroid therapy in the treatment of any process, condition, event or disease as hereinbefore set forth; or
  - c) for co-administration together with a glucocorticosteroid in the treatment of any process, condition, event or disease as hereinbefore set forth; and/or.
- V. A pharmaceutical dosage form comprising a 5-HT<sub>3</sub> receptor antagonist together with a glucocorticosteroid, especially for the local treatment of any process, condition, event or disease as hereinbefore set forth.

Where under (I) to (V) the term "any process, condition, event or disease" is used, this term preferably relates to the diseases mentioned under (1), (2) and (3) above, especially as defined above as being preferred.

The term "locally administering" or "local use" is defined to mean local administration or especially "local treatment" as defined above, that is, administration at or near the site of the non-inflammatory disease, irritation condition or local inflammatory disease condition, in contrast to systemic administration.

Dosage forms, e.g. in accordance with V above, are to be understood as including both fixed-unit-dosage forms, e.g. liquid formulations, comprising both active ingredients together

with appropriate pharmaceutically acceptable diluents or carriers, as well as twin delivery systems, packages or the like comprising both active ingredients separately or in separate dosage form, for concomitant or sequential administration.

The 5-HT<sub>3</sub> receptor antagonists are preferably used in well-known liquid formulations.

Utility of 5-HT<sub>3</sub> receptor antagonists in accordance with the present invention can be demonstrated in clinical trials carried out in accordance with standard techniques and methodologies, for example as follows:

The following examples are for illustrative purposes and are not intended to diminish the scope of the present invention. Instead of tropisetron, any other 5-HT<sub>3</sub>-antagonist, or a pharmaceutically acceptable salt thereof, solvate, e.g. hydrate, or crystalline form thereof, especially selected from the group consisting of ondansetron, granisetron, dolasetron, ramosetron, fabesetron, lintopride and alosetron, can be used, or any combination of two or more of these 5-HT<sub>3</sub> receptor antagonists or pharmaceutically acceptable salts thereof.

In the following examples, tropisetron is administered in the standard formulation of the trademark Navoban® which is available in ampoules that contain 2 mg or 5 mg of the active substance, tropisetron.

EXAMPLE 1: Treatment of synovial inflammation/synovitis consequent to inflammatory processes

Trials are performed on 2 patients exhibiting rheumatoid arthritis and severe consequential synovial inflammation as well as marked pain.

Two patients exhibiting acute exacerbation of rheumatoid arthritis, one in the shoulder joint, the other in the knee joint, are each treated once with 2 mg tropisetron administered intra-articularly. In both cases treatment leads to almost complete remission from pain within a few hours. Both patients are examined over a period of one week following treatment with tropisetron and are found to be free of symptoms.

**EXAMPLE 2: Treatment of synovial inflammation following traumatic or degenerative event**

A first patient exhibits repeated exudation from the knee joint consequential to damage to the meniscus. Prior to the trial the patient received injections of glucocorticoid. The patient receives an injection of 2 mg of tropisetron administered intra-articularly. After 45 min. marked reduction of pain is reported and a major reduction of effusion from the knee joint is observed after 5 hours. The patient remains free of symptoms without further therapy over an observation period of 5 days.

A second patient exhibits damage to the meniscus of the right knee joint as well as arthritic change leading to synovial irritation with consequential knee joint effusion. Despite a successful synovectomy prior to trial entry the patient exhibits renewed knee joint effusion. Two 2 mg doses of tropisetron are administered i.v. over a period of 17 days with 15 injections, one each day with a 2 days pause in therapy. 24 hours after the first i.v. injection, significant improvement of pain is reported. Following continuation of injections, the patient exhibits as virtually free of symptoms. The exudation from the knee joint is completely inhibited without any other medication within 8 days and movement of the knee joint is clearly improved. The improvement in condition continues over a further 7 days observation following completion of therapy:

**EXAMPLE 3: Treatment of implant or insertion tendopathy**

Three patients exhibiting implant/insertion tendopathy are treated with tropisetron. The condition treated results in the case of the first patient from epicondylitis radialis, in the case of the second patient from insertion tendopathy in the area of insertion of the deltoid muscle and, in the case of the third patient from an enthesiopathy of the Malleolus lateralis. In all three cases, tropisetron is injected once at a dosage of 2 mg in the area of greatest pain, directly into the tissues. In all three cases severity of pain decreases within a period of 60 min. and virtually complete remittal is achieved within 24 hrs., indicating remission from disease remission. Further improvements found after tropisetron injection include increase of strength and of flexibility of the affected body part.

**EXAMPLE 4: Further Examples for the efficiency of tropisetron:**

Pain is subsequently determined on a visual analogue scale (VAS) that goes from 0 mm (no pain) to 100 mm (strongest pain).

4 a) Enthesiopathy (Insertion Tendopathy) (with infiltration):

- (i) Female patient, born in 1955; Diagnosis: epicondylitis. Rest pain before local injection of tropisetron 30 mm, 7 days later 0 mm.
- (ii) Female patient, born 1947, diagnosis: periarthropathia humeroscapularis. Rest pain before local injection of tropisetron 85 mm, 7 days after local injection 45 mm.
- (iii) Patient born 1953, diagnosis: epicondylitis. Rest pain before local injection of tropisetron 35 mm, 7 days thereafter 0 mm.

4 b) Rheumatoid arthritis (intraarticular injection):

- (i) Female patient, born 1942.

Before injection of Navoban into the right knee joint: rest pain 62 mm, pain under exercise 82 mm, bending up to 105 °, knee joint circumference 46 cm. 24 hours later: rest pain 22 mm, pain under exercise 21 mm, bending 105 °, knee joint circumference 45 cm. 48 hours after treatment: rest pain 0, pain under exercise 5 mm, bending 110 °, knee joint circumference 45 cm. 7 days after treatment: rest pain 20 mm, pain under exercise 16 mm, bending up to 145 °, knee joint circumference 45.4 cm.

- (ii) Female patient, born 1947.

Before injection of 2 mg Navoban into the left knee joint: rest pain 25 mm, pain under exercise 85 mm, bending up to 115 °, knee joint circumference 37.5 cm. 24 hours later: rest pain 8 mm, pain under exercise 58 mm, bending 120 °, knee joint circumference 37.5 cm. 48 hours after treatment: rest pain 6 mm, pain under exercise 55 mm, bending 115 °, knee joint circumference 37.5 cm. 7 days after treatment: rest pain 5 mm, pain under exercise 59 mm, bending up to 115 °, knee joint circumference 37 cm. Now 5 mg Navoban are injected into the knee joint, 7 days later the patient has a rest pain of 0 mm, pain under exercise of 33 mm, bending is possible up to 120 °, and the knee joint circumference is 37.5 mm.

4 c) Treatment of left knee joint irritation (Morbus Still):

Patient, 26 year old, Morbus Still diagnosed 1.5 years ago. Despite basic treatment with 15 mg methotrexat per week and 20 mg prednisolone per day, the patient suffers from swelling

and overheating of the left knee joint for the first time in February 1999. Pain at rest in the VAS 98 (0-100), pain under strain 70. Obvious articular effusion. Knee joint circumference 44 cm, straightening/bending 0°-10°, 115°. 24 hours after an intraarticular injection with 2 mg tropisetron pain at rest in the Visual Analog Scale (VAS) 0, pain under strain 70. Straightening/bending 0°, 0°, 135°, knee joint circumference 41 cm, with significant reduction of joint swelling and effusion respectively. 7 days later pain at rest in the VAS 0, pain under strain 57, straightening/bending 0°, 0°, 130°, knee joint circumference 40.5 cm.

#### 4 e) Gonarthrosis:

Patient, 74 years old, has activated gonarthrosis with effusion for 2 months. Only short-term pain reduction by means of non-steroidal antiphlogistics could be achieved. Following an intra-articular injection with 2 mg tropisetron, pain at rest after 24 hours on the visual analog scale for pain (0-100) sinks from 43 to 25, after 2 days to 13, after 7 days to 10 and in a follow-up check after 17 days to 4. The analog values for pain under strain are 75, 45, 10 and 10, respectively. The effusion is gone after 7 days, as is the overheating of the knee joint.

#### Example 5: Spinal syndrome (i.v. administration)

Female patient born 1957, diagnosis: cervical syndrome. Before injection of 2 mg tropisetron 68 mm rest pain, 24 hours later 35 mm, at day 7 18 mm.

#### Example 6: Spinal syndromes (i.v. administration)

##### (i) Cervical Syndrome:

Female patient born 1957, diagnosis: cervical syndrome. Before i.v. injection of 2 mg tropisetron 68 mm rest pain, 24 hours later 35 mm, at day 7 18 mm.

##### (ii) Lumbalgia:

Patient born 1940, diagnosis: lumbalgia. The patient is injected 2 mg Navoban i.v., the rest pain before the injection amounts to 78 mm, the pain under exercise 90 mm, the Schober distance upright/with bended upper part of the body between the processus spinalis vertebrae of vertebra 1 and 5 ("Schober" hereinafter) 10/11 cm, the distance from fingers to ground with bended upper part of the body 32 cm; 24 hours later, the rest pain is 36 mm, the pain under exercise 43 mm, the Schober 10/12.3, and the finger/ground distance 24 cm; after 48 hours, the rest pain is 28 mm, the pain under exercise 47 mm, the Schober

10/12.5, the finger/ground distance 26.5 cm. 7 days after treatment, the rest pain is 15 mm, the pain under exercise 22 mm, the Schober 10/13, and the finger/ground distance 22 cm.

Equivalent results as in the preceding examples are obtainable in equivalent or comparable trials with patients exhibiting similar symptomatology employing 5-HT<sub>3</sub>-receptor antagonists other than tropisetron, for example using any of the 5-HT<sub>3</sub>-receptor antagonists A) through C) or E) through H) hereinbefore recited at comparable, e.g. conventional clinical, dose as known in the art. Similar results are also achievable employing 5-HT<sub>3</sub>-receptor antagonists, e.g. tropisetron at doses of the order of 2mg/day p.o. or by injection or topical application in clinical trials involving subjects exhibiting other local non-inflammatory or local inflammatory diseases, conditions or symptoms.

Trials conducted as described above or analogously are demonstrative of long lasting and disease modifying effects in conditions herein described as well as symptomatic and glucocorticosteroid replacement effect for 5-HT<sub>3</sub> receptor antagonists.

For use in accordance with the present invention the appropriate dosage will, of course, vary depending on for example the particular 5-HT<sub>3</sub> receptor antagonist employed the mode of administration and the nature and severity of the condition to be treated as well as the specific condition to be treated. In general an indicated single, e.g. daily, dosage will be in the range usually employed for known indications such as emesis and will typically be from about 0.05 to about 50 mg per day, more preferably around 1 to 10 mg per day, conveniently administered once or in divided doses up to four times a day or in sustained release form, or used repeatedly after longer intervals, e.g. after some days or weeks, e.g. after 2 days to 4 weeks. In the case of tropisetron an appropriate dosage for administration, e.g. by injection, for example for i.v. application or injection direct into the affected areas, will be of the order of 2 mg per day or 5 mg per day, administered once, sequentially over a sequence of 2 to 20 days or at intervals of 2 to 5 days to 2 days to 2 weeks.

For use in accordance with the invention, 5-HT<sub>3</sub> receptor antagonists may be administered by any conventional route in particular enterally, preferably orally, e.g. in the form of tablets or capsules, or rectally, e.g. in the form of suppositories or enema, or most preferably parenterally, e.g. in the form of injectible solutions or suspensions, e.g. by subcutaneous, intraperitoneal or intra-muscular injection for systemic administration. Suitable formulations

for use in accordance with the present invention will include any of those as known and commercially available and clinically employed in the art, for example the commercially available formulations. Preferably, the compositions are administered locally, that is, near or at the site of the manifestation of the disease to be treated, e.g. by intramuscular injection, intra-articular injection or any other injection near or at the site of disease manifestation; in a broader aspect of the invention, also local tissue infiltration or transdermal administration may be considered, e.g. by use of gels, creams or ointments or the like, or preferably by transdermal patches. Dosages for such forms will be of the order or slightly higher than those used on administration by injection.



CLAIMS

1. The use of a 5-HT<sub>3</sub> receptor antagonist or of a pharmaceutically acceptable salt of such an antagonist for the manufacture of a pharmaceutical composition for the treatment of a non-inflammatory local disease of the musculo-skeletal system, of a local irritation condition of a joint or tendon sheath; or for the local treatment a local manifestation at the locomotor apparatus of an inflammatory disease except for a crystal induced arthritis and a living pathogen induced inflammatory disease condition as long as the living pathogen is still present.
2. The use according to claim 1, wherein in addition to pain at least one further sequela or symptom of the local disease, local irritation condition or local manifestation is alleviated, ameliorated or controlled.
3. The use according to claim 1, wherein the 5-HT<sub>3</sub> receptor antagonist is selected from the group consisting of ondansetron, granisetron, dolasetron, tropisetron, ramosetron, fabesetron, lintopride and alosetron, which may be used in free form or as a pharmaceutically acceptable salt.
4. The use according to claim 1, wherein the disease to be treated is a non-inflammatory local disease of the musculo-skeletal system selected from the group consisting of myofasciale syndrome, tendomyosis, tendinosis, insertion tendopathy, bursopathy or periarthopathy, overload syndrome of the muscle, syndromes due to the compression of nerves or neuropathy and algodystrophy.
5. The use according to claim 1, wherein the disease to be treated is a non-inflammatory local disease of the musculo-skeletal system selected from the group consisting of myofasciale syndrome, tendomyosis, tendinosis, insertion tendopathy, bursopathy or periarthopathy, overload syndrome of the muscle, and syndromes due to the compression of nerves.
6. The use according to claim 1, wherein the disease to be treated is a local irritation condition of a joint or tendon sheath related to a meniscus lesion, arthrosis, trauma,

locally osteochondritis dissecans, osteonecrosis or joint chondromatosis, and various non-inflammatory rheumatoid diseases.

7. The use according to claim 1 for the local treatment a local manifestation at the locomotor apparatus of an inflammatory disease except for a crystal induced arthritis and a living pathogen induced inflammatory disease condition as long as the living pathogen is still present, wherein said inflammatory disease being selected from chronic polyarthritis, sarcoidosis, paraneoplastic syndrome or tumor-induced inflammatory diseases, turbid effusions, collagenosis, postinfectious arthritis, seronegative spondylarthritis and vasculitis.
8. A method of treating a non-inflammatory local disease of the musculo-skeletal system, of treating a local irritation condition of a joint or tendon sheath, or of locally treating a local manifestation at the locomotor apparatus of an inflammatory disease except for a crystal induced arthritis and a living pathogen induced inflammatory disease condition as long as the living pathogen is still present, in a subject in need thereof, which method comprises locally administering to said subject an effective amount of a 5-HT<sub>3</sub> receptor antagonist.
9. A method according to claim 8 wherein the 5-HT<sub>3</sub> antagonist is administrated locally at or near the site of the local disease, local irritation condition or local manifestation.
10. A method of providing replacement therapy for glucocorticosteroid therapy in a subject receiving such glucocorticosteroid therapy, which method comprises locally administering to said subject an effective amount of a 5-HT<sub>3</sub>-receptor antagonist.
11. A method of treating any process, condition, event or disease as hereinbefore described, in a subject in need thereof, which method comprises locally administering an effective amount of a 5-HT<sub>3</sub> receptor antagonist together with a glucocorticosteroid.
12. A pharmaceutical dosage form for local administration comprising a 5-HT<sub>3</sub> receptor antagonist together with an anti-inflammatory glucocorticosteroid.

13. A 5-HT<sub>3</sub> receptor antagonist or a pharmaceutical composition comprising a 5-HT<sub>3</sub> receptor antagonist for local use:
- a) in the treatment of process, condition, event or disease as hereinbefore described;
  - b) as replacement therapy for anti-inflammatory glucocorticosteroid therapy in the treatment of any process, condition, event or disease as hereinbefore described; or
  - c) for co-administration together with a glucocorticosteroid in the treatment of any process, condition, event or disease as hereinbefore described.
14. A method, use, compound or composition as claimed in any one of claims 1, 8, 10, 11, 12 or 13, wherein the 5-HT<sub>3</sub> receptor antagonist is tropisetron.